

**ASAM Patient Placement Criteria 2-Revised
Adult Risk Rating → Recommended Level of Care Matrix**

Dimension	Appropriate Levels of Care by Risk Rating					
	0	1	2	3	4 or 4a	4b
Dim. One	No Tx Recommended	Level I-D	Level II-D	Level III.2-D Level III.7-D	Level IV-D	-
Dim. Two	No Tx Recommended	Level I	Level I Level II.1	Level III.7	Level IV	-
Dim. Three	No Tx Recommended	Level I	Level II.1 Level II.5	Level III.5 Level III.7	Level IV	-
Dim. Four	No Tx Recommended	Level I	Level I Level II.1	Level I Level II.1	Level I Level II.1	Level III.3 Level III.7 Level IV
Dim. Five	No Tx Recommended	Level I	Level II.1 Level II.5	Level II.5 Level III.1	Level I Level II.1 Level III.1 Level III.5	Level III.3 Level III.5 Level IV
Dim. Six	No Tx Recommended	Level I	Level I Level II.1 Level II.5 Level III.1	Level II.1 Level II.5 Level III.1	Level III.1	Level III.1 Level III.3 Level III.5

Always refer to the PPC manual (Appendix A) for more detailed information and rationales for these recommendations

The highest dimensional LOC recommendation should determine the actual initial placement. Patient placement must be reviewed periodically using PPC continued stay / discharge criteria. At a minimum, this must occur during treatment plan reviews.

No Tx Rec.: Referral to Self-Help Groups or Level 0.5
Early Intervention / Education may be appropriate.
Level III.2-D = Sub-Acute Detoxification
Level III.7-D / IV-D = Acute Detoxification

Level I = Outpatient
Level II.1 = Intensive Outpatient
Level II.5 = IOP Day Treatment
Level III.1 = Recovery House

Level III.3 = Long-Term Residential
Level III.5 = Clinically Managed Intensive Inpatient
Level III.7 = Medically Monitored Intensive Inpatient
Level IV = Medically Managed Intensive Inpatient