

Eligible Conditions With Age and Type of Evidence

All disability conditions must originate prior to age 18, be expected to continue indefinitely, and result in substantial limitation in adaptive functioning per WAC 388-823.

Eligible Conditions Specific to Age				
Condition	0 to 3	4 to 9	10 to 17	18 and older
Developmental Delays	X	X		
Intellectual Disability (ID)		X	X	X
Cerebral Palsy		X	X	X
Epilepsy		X	X	X
Autism		X	X	X
Another neurological or other condition similar to Intellectual Disability		X	X	X

Tests Accepted as Evidence of Substantial Limitation	
Cognitive Tests	Adaptive Assessments
<ul style="list-style-type: none"> Stanford-Binet Wechsler Intelligence Scales Differential Abilities Scale (DAS) Kaufman Assessment Battery for Children (K-ABC) Das-Naglieri Cognitive Assessment System (CAS) Woodcock Johnson Test of Cognitive Abilities (WJ) <p>If you have a hearing impairment, English is not your primary language or you are non-verbal, your FSIQ may be estimated using one of the tests below:</p> <ul style="list-style-type: none"> Leiter International Performance Scale-Revised (Leiter-R) Wechsler Intelligence Scales Performance scale Comprehensive Test of Nonverbal Intelligence (C-TONI) Kaufman Assessment Battery for Children (K-ABC) Nonverbal scale index 	<ul style="list-style-type: none"> Vineland Adaptive Behavior Scales (VABS) Scales of Independent Behavior-Revised (SIB-R) Inventory for Client and Agency Planning (ICAP) Adaptive Behavior Assessment System (ABAS)

Diagnosis, Diagnostician, and Required Evidence of Substantial Limitation		
Diagnosis	Diagnostician	Substantial Limitation
Developmental Delay	Not applicable	Developmental Delays
Intellectual Disability	Licensed Psychologist or Certified School Psychologist	Adaptive functioning assessment with a standard score of ≤ 69 , and Psychological assessment with FSIQ of ≤ 69 or Stanford-Binet IV with FSIQ of ≤ 67
Cerebral Palsy, or Similar brain damage which causes, quadriplegia, hemiplegia, or diplegia	Licensed Physician	Onset prior to age 3, and Evidence of the need for direct physical assistance in any 2 of the following: toileting, bathing, eating, dressing, mobility or communication
Epilepsy, Seizure Disorder	Board Certified Neurologist	Seizures are uncontrolled and ongoing or recurring, and Adaptive functioning assessment with a standard score of ≤ 69
Autism, Autistic Disorder (DSM-IV-TR-299.00), or Autism Spectrum disorder (DSM-5) with severity level of 2 or 3 in both columns	Board Certified Neurologist; Board Certified Psychiatrist; Licensed Psychologist; Board Certified Developmental and Behavioral Pediatrician; Licensed Physician or ARNP associated with an Autism, Developmental Center, or Center of Excellence	Onset prior to age 3, and Adaptive functioning assessment with a standard score of ≤ 69 , and For diagnosis from DSM-5 FSIQ of ≤ 84 , or evidence that person meets DSM-IV-TR criteria
Another neurological or other condition similar to Intellectual Disability	Licensed Physician	Adaptive functioning assessment with a standard score of ≤ 69 , and FSIQ of ≤ 77 or Stanford-Binet IV with FSIQ of ≤ 75 , or if under age 20, scores in both Broad Reading & Broad Math ≤ 69 can replace FSIQ testing